

# Abacavir sulfate, Lamivudine, and Zidovudine



**Brand Name:** Trizivir

## Drug Description

Abacavir/lamivudine/zidovudine (for brevity, from this point referred to as Trizivir) is a fixed-dose tablet containing three synthetic nucleoside analogues: abacavir sulfate, lamivudine, and zidovudine. Each tablet contains 300 mg of abacavir sulfate, 150 mg of lamivudine, and 300 mg of zidovudine, each of which inhibits HIV-1 viral reverse transcriptase. [1]

## HIV/AIDS-Related Uses

Trizivir was approved by the FDA on November 14, 2000, for use in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults or adolescents weighing more than 40 kg.[2] Based on interim results from a current clinical trial, therapy with Trizivir alone is not recommended. The interim report indicates that treatment-naïve patients given Trizivir alone experienced virologic failure earlier and more frequently than those given Trizivir in combination with other antiretrovirals.[3] [4]

## Pharmacology

Each of the three synthetic nucleoside analogues contained in Trizivir inhibits viral reverse transcriptase (RT), an enzyme HIV requires in order to replicate. Abacavir, lamivudine, and zidovudine work by incorporating themselves into viral DNA and terminating the viral DNA chain. (For more information, see individual drug fact sheets for abacavir sulfate, lamivudine, and zidovudine.)

Abacavir is a carbocyclic nucleoside analogue that is converted by cellular enzymes to the active metabolite, carbovir triphosphate. Carbovir triphosphate is an analogue of deoxyguanosine-5'-triphosphate (dGTP). Carbovir triphosphate inhibits RT by competing with the natural substrate dGTP and by incorporation into viral DNA. The lack of a 3'-OH group in the incorporated nucleoside analogue prevents the formation of the 5' to 3' phosphodiester linkage essential for DNA chain elongation. In vitro, abacavir had synergistic activity in combination

with amprenavir, nevirapine, and zidovudine, and additive activity with didanosine, lamivudine, stavudine, and zalcitabine. Following oral dosing, abacavir is rapidly absorbed and extensively distributed. Binding of abacavir to human plasma proteins is about 50%, independent of concentration. Abacavir is primarily eliminated by metabolism by alcohol dehydrogenase to form the 5'-carboxylic acid and glucuronyl transferase to form the 5'-glucuronide.[5]

Lamivudine is a synthetic nucleoside analogue that is phosphorylated intracellularly to its active 5'-triphosphate metabolite, lamivudine triphosphate (L-TP). L-TP inhibits viral RT by DNA chain termination. In vitro, lamivudine had synergistic antiretroviral activity with zidovudine. Following oral dosing, lamivudine is rapidly absorbed and extensively distributed. Plasma protein binding is low and about 70% of an intravenous dose is excreted unchanged in the urine. Metabolism is a minor route of elimination.[6]

Zidovudine is phosphorylated intracellularly to its active 5'-triphosphate metabolite, zidovudine triphosphate (ZDV-TP). ZDV-TP also inhibits RT by DNA chain termination. In vitro, zidovudine demonstrates synergistic activity with delavirdine, didanosine, indinavir, nelfinavir, nevirapine, ritonavir, saquinavir, and zalcitabine, and additive activity with interferon-alpha. Following oral dosing, zidovudine is rapidly absorbed and extensively distributed. Plasma protein binding is low and elimination is primarily by hepatic metabolism. The major metabolite is 3'-azido-3'-deoxy-5'-O-beta-D-glucopyranuronosylthymidine. A second metabolite, 3'-amino-3'-deoxythymidine, has been identified.[7]

In a bioavailability study of Trizivir compared to separate tablets of the three components given simultaneously to healthy adults, there was no difference in absorption. One Trizivir tablet was bioequivalent to dosing with one tablet each of abacavir sulfate 300 mg, lamivudine 150 mg, and zidovudine 300 mg in healthy, fasting adults.[8]

Trizivir is in FDA Pregnancy Category C. No

# Abacavir sulfate, Lamivudine, and Zidovudine



## Pharmacology (cont.)

---

adequate or well-controlled studies of the combination drug have been done in pregnant women. A study of zidovudine therapy in women in the last trimester of pregnancy showed that although this drug does cross the placenta, there was no evidence of drug accumulation, and zidovudine concentrations in neonatal plasma at birth were essentially equal to those in maternal plasma at delivery. Studies in laboratory animals have shown that abacavir and lamivudine cross the placenta, with evidence of fetal toxicity at dosage levels many times higher than the corresponding dose for humans. Trizivir should be used in pregnancy only if the potential benefits outweigh the risks. An Antiretroviral Pregnancy Registry has been established to monitor the outcomes of pregnant women exposed to Trizivir and other antiretrovirals. Physicians may register patients by calling (800) 258-4263. Zidovudine is excreted in human milk, and abacavir and lamivudine are excreted in the milk of laboratory animals.[9]

HIV-1 isolates with reduced sensitivity to abacavir, lamivudine, or zidovudine have been selected in vitro and have also been obtained from patients treated with that combination or lamivudine plus zidovudine. Treatment for 12 weeks with lamivudine and zidovudine restored sensitivity to zidovudine in some patients with zidovudine-resistant virus. Combination therapy delayed the emergence of mutations conferring resistance to zidovudine. Higher levels of resistance were associated with greater numbers of mutations. Laboratory strains of HIV-1 containing multiple RT mutations conferring abacavir resistance exhibited cross-resistance to lamivudine, didanosine, and zalcitabine in vitro. Cross-resistance to didanosine and zalcitabine has been observed in some patients who harbor lamivudine-resistant HIV-1 isolates. Multiple drug resistance, including resistance to lamivudine and stavudine, has been observed in HIV isolates from patients treated for more than 1 year with zidovudine plus didanosine or zalcitabine.[10]

Interim results from a current study show that antiretroviral-naïve patients randomized to receive Trizivir alone experienced virologic failure earlier and more frequently than patients who were

randomized to receive either of the two other regimens being evaluated in the study. The two other regimens were: (1) a combination of lamivudine/zidovudine plus efavirenz, and (2) the combination of Trizivir with efavirenz. Virologic failure was defined as having an HIV RNA level in plasma above 200 copies/ml at least 4 months after starting study treatment. After an average of 32 weeks on study, 21% in the group receiving Trizivir experienced virologic failure versus 10% in the other two groups combined. Virologic failure occurred sooner and more often in those receiving Trizivir alone, regardless of their initial viral load (whether above or below 100,000 copies/mL).[11]

## Adverse Events/Toxicity

---

Trizivir contains abacavir sulfate, which has been associated with fatal hypersensitivity reactions. In clinical studies, approximately 5% of adult and pediatric patients receiving abacavir developed a hypersensitivity reaction. Hypersensitivity reactions are characterized by symptoms indicating multi-organ/body system involvement, usually appearing within the first 6 weeks of abacavir therapy, although they may appear at any time. Frequently observed signs and symptoms of hypersensitivity include fever, skin rash, fatigue, and gastrointestinal symptoms such as nausea, vomiting, diarrhea, or abdominal pain. Other signs and symptoms include malaise, lethargy, myalgia, myolysis, arthralgia, edema, cough, dyspnea, headache, and paresthesia. The diagnosis of hypersensitivity reaction should be considered for patients presenting with symptoms of acute onset respiratory diseases such as pneumonia, bronchitis, or flu-like illnesses. Physical findings associated with hypersensitivity reactions include lymphadenopathy, mucous membrane lesions (conjunctivitis and mouth ulcerations), and sometimes a maculopapular or urticarial rash. Laboratory abnormalities include elevated liver function tests, increased creatine phosphokinase or creatinine, and lymphopenia. Anaphylaxis, liver failure, renal failure, hypotension, and death have occurred in association with hypersensitivity reactions. Severe or fatal hypersensitivity reactions can occur within hours after reintroduction of abacavir in patients who have no identified history or who have unrecognized symptoms of

# Abacavir sulfate, Lamivudine, and Zidovudine



## Adverse Events/Toxicity (cont.)

---

hypersensitivity to abacavir. Trizivir should be discontinued permanently if hypersensitivity cannot be ruled out.[12]

Lactic acidosis and severe hepatomegaly with steatosis have been reported with the use of nucleoside analogues alone or in combination. These conditions are sometimes fatal. Female gender, obesity, and prolonged nucleoside exposure may be risk factors. Caution should be exercised in any patient with known risk factors for liver disease; however, liver problems have been reported in patients with no known risk factors. Treatment with Trizivir should be suspended in any patient who develops clinical or laboratory findings that suggest the presence of lactic acidosis or pronounced hepatotoxicity.[13] Granulocytopenia and anemia are the most frequent adverse effects associated with zidovudine therapy.[14] Myopathy and myositis have occurred with prolonged use of zidovudine and may occur during therapy with Trizivir.[15] Peripheral neuropathy has been reported in adults receiving lamivudine but has rarely resulted in interruption or discontinuance of treatment.[16] Post-treatment exacerbations of hepatitis B virus (HBV) infections have been reported in both HIV and non-HIV infected patients treated with lamivudine for chronic HBV when lamivudine therapy was discontinued.[17]

Other adverse effects occurring in clinical trials of Trizivir or its component drugs include nausea, vomiting, diarrhea, abdominal pain or cramping, dyspepsia, anorexia, insomnia and other sleep disorders, fever and/or chills, headache, dizziness, malaise and/or fatigue, depressive disorders, neuropathy, musculoskeletal pain, myalgia, arthralgia, and skin rashes. Adverse events reported during post-approval use of abacavir, lamivudine, and/or zidovudine that may potentially be related to these drugs include cardiomyopathy, stomatitis, oral mucosal pigmentation, gynecomastia, hyperglycemia, vasculitis, weakness, anemia, aplastic anemia, lymphadenopathy, pure red cell aplasia, splenomegaly, lactic acidosis and hepatic steatosis, pancreatitis, post-treatment exacerbation of hepatitis B, sensitization reactions and urticaria, muscle weakness, CPK elevation, rhabdomyolysis, paresthesia, peripheral neuropathy, seizures,

abnormal breath sounds/wheezing, alopecia, erythema multiforme, and Stevens-Johnson syndrome.[18]

## Drug and Food Interactions

---

Trizivir may be administered with or without food. Administration of Trizivir with food did not alter the extent of abacavir, lamivudine, and zidovudine absorption (AUC), as compared to administration under fasting conditions.[19]

Abacavir, lamivudine, and zidovudine (studied as individual drugs) are not significantly metabolized by the cytochrome P450 enzymes; therefore, it is unlikely that clinically significant drug interactions will occur with drugs metabolized through these pathways.[20]

Abacavir administered at twice the recommended dose increased methadone clearance by 22%. A small number of patients receiving both abacavir and methadone may need a methadone dosage adjustment. Because abacavir elimination is decreased by alcohol, consumption of alcohol may cause an increase in abacavir exposure.[21]

Because lamivudine and zalcitabine may inhibit the intracellular phosphorylation of one another, Trizivir should not be co-administered with zalcitabine.[22] Lamivudine exposure (AUC) was increased by 44% and lamivudine renal clearance was decreased by 30% when co-administered with sulfamethoxazole/trimethoprim. Concurrent administration of lamivudine and zidovudine in one small study resulted in a 39% increase in peak plasma concentration of zidovudine with no change observed in AUC. Concurrent administration of lamivudine with indinavir and zidovudine resulted in a 6% decrease in AUC of lamivudine, no change in AUC of indinavir, and a 36% increase in AUC of zidovudine. No adjustment in dose is necessary. Concurrent administration of lamivudine with drugs associated with pancreatitis (alcohol, didanosine, IV pentamidine, sulfonamides, and zalcitabine) or with drugs associated with peripheral neuropathy (dapsone, didanosine, isoniazid, stavudine, and zalcitabine) should be avoided or used cautiously.[23]

Zidovudine may interact with atovaquone,



# Abacavir sulfate, Lamivudine, and Zidovudine



## Drug and Food Interactions (cont.)

clarithromycin, fluconazole, methadone, phenytoin, probenecid, ribavirin, rifampin, valproic acid,[24] nelfinavir, and ritonavir.[25] The hematologic toxicity of zidovudine may be increased when zidovudine is co-administered with bone marrow depressant agents such as ganciclovir or interferon-alpha and others, or blood dyscrasia-causing medications or cytotoxic agents, as well as with radiation therapy.[26] [27] Medications that are metabolized by hepatic glucuronidation such as acetaminophen, aspirin, benzodiazepines, cimetidine, indomethacin, morphine, and sulfonamides may in theory increase the risk of toxicity of zidovudine or the other medication.[28] An antagonistic relationship between zidovudine and stavudine, doxorubicin, and ribavirin has been reported in vitro. Concomitant use of zidovudine with any of these three drugs should be avoided.[29]

## Contraindications

Trizivir tablets are contraindicated in patients with previously demonstrated hypersensitivity to any of the components of the product. Trizivir contains abacavir sulfate, which has been associated with fatal hypersensitivity reactions. An actual or suspected hypersensitivity reaction to abacavir sulfate is an absolute contraindication to Trizivir use.

Due to the fixed-dose formulation of Trizivir, there is no way to accommodate the dosage reduction of zidovudine that may be necessary in individuals with impaired liver function or the dosage adjustment of both lamivudine and zidovudine that may be necessary in those with renal insufficiency (creatinine clearance less than 50 ml/min). Additionally, dosage adjustments cannot be made for pediatric or geriatric patients or for patients who weigh less than 40 kg or for any patient with special dosing requirements. Trizivir is not recommended for these patients.[30]

## Clinical Trials

For information on clinical trials that involve Abacavir sulfate, Lamivudine, and Zidovudine, visit the ClinicalTrials.gov web site at

<http://www.clinicaltrials.gov>. In the Search box, enter: Abacavir sulfate, Lamivudine, and Zidovudine AND HIV Infections.

## Dosing Information

Mode of Delivery: Oral.[31]

Dosage Form: Oral film-coated tablet. Each tablet contains 300 mg of abacavir sulfate, 150 mg of lamivudine, and 300 mg of zidovudine. The recommended dose for adults and adolescents is 1 tablet twice daily. Because this medication is in a fixed-dose tablet, it is not recommended for use in adults or adolescents who weigh less than 40 kg (88 lbs) or in patients requiring dose adjustment such as those with creatinine clearance less than 50 ml/min or those experiencing dose-limiting adverse events.[32]

Storage: Store at 25 C (77 F); excursions permitted to 15 C to 30 C (59 F to 86 F).[33]

## Chemistry

CAS Name: Abacavir sulfate: (1S,4R)-4-[2-Amino-6-(cyclopropylamino)-9H-purin-9-yl]-2-cyclopentene-1-methanol sulfate[34]

Lamivudine: 2(1H)-Pyrimidinone, 4-amino-1-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]-, (2R-cis)-[35]

Zidovudine: Thymidine, 3'-azido-3'-deoxy-[36]

CAS Number: Abacavir sulfate: 188062-50-2[37]

Lamivudine: 134678-17-4[38]

Zidovudine: 30516-87-1[39]

Molecular formula: Abacavir sulfate: C14-H18-N6-O<sub>11</sub>/2H<sub>2</sub>O-4-S; Lamivudine: C<sub>8</sub>-H<sub>11</sub>-N<sub>3</sub>-O<sub>3</sub>-S; Zidovudine: C<sub>10</sub>-H<sub>13</sub>-N<sub>5</sub>-O<sub>4</sub>[40]

Abacavir sulfate: C50.1%, H5.7%, N25.1%, O14.3%, S4.8%]; Lamivudine: C41.91%, H4.84%, N18.33%, O20.94%, S13.99%; Zidovudine: C44.94%, H4.90%, N26.21%, O23.95%[41]

# Abacavir sulfate, Lamivudine, and Zidovudine



## Chemistry (cont.)

---

Molecular weight: Abacavir sulfate: 670.76;  
Lamivudine: 229.26; Zidovudine: 267.24[42]

Melting point: Abacavir: 165 C; Lamivudine:  
160-162 C; Zidovudine: 106-112 C[43]

Physical Description: Film-coated tablets  
containing 300 mg of abacavir as abacavir sulfate,  
150 mg of lamivudine, and 300 mg of zidovudine  
plus inactive ingredients.[44]

Solubility: Abacavir sulfate: 77 mg/ml in distilled  
water at 25 C; Lamivudine: 70 mg/ml in water at 20  
C; Zidovudine: 20.1 mg/ml in water at 25 C.[45]

## Further Reading

---

Fischl MA, Burnside AE Jr, Farthing CE,  
Thompson MA, Bellos NC, Williams VC, Kauf  
TL, Wannamaker PG, Shaefer MS; ESS40005  
Study Team. Twice-daily Trizivir versus  
Combivir-abacavir in antiretroviral-experienced  
adults with human immunodeficiency virus-1  
infection: a formulation-switch trial.  
Pharmacotherapy. 2003 Nov;23(11):1432-40.

Katlama C, Fenske S, Gazzard B, Lazzarin A,  
Clumeck N, Mallolas J, Lafeuillade A, Mamet JP,  
Beauvais L; AZL30002 European study team.  
TRIZAL study: switching from successful HAART  
to Trizivir (abacavir-lamivudine-zidovudine  
combination tablet): 48 weeks efficacy, safety and  
adherence results. HIV Med. 2003 Apr;4(2):79-86.

Loneragan JT, Barber RE, Mathews WC. Safety and  
efficacy of switching to alternative nucleoside  
analogues following symptomatic hyperlactatemia  
and lactic acidosis. AIDS. 2003 Nov  
21;17(17):2495-9.

John M, McKinnon EJ, James IR, Nolan DA,  
Herrmann SE, Moore CB, White AJ, Mallal SA.  
Randomized, controlled, 48-week study of  
switching stavudine and/or protease inhibitors to  
combivir/abacavir to prevent or reverse lipodystrophy  
in HIV-infected patients. J Acquir Immune Defic  
Syndr. 2003 May 1;33(1):29-33.

Ruane PJ, Parenti DM, Margolis DM, Shepp DH,  
Babinchak TJ, Van Kempen AS, Kauf TL,

Danehower SA, Yau L, Hessenthaler SM, Goodwin  
D, Hernandez JE; COL30336 Study Team.  
Compact quadruple therapy with the  
lamivudine/zidovudine combination tablet plus  
abacavir and efavirenz, followed by the  
lamivudine/zidovudine/abacavir triple nucleoside  
tablet plus efavirenz in treatment-naïve  
HIV-infected adults. HIV Clin Trials. 2003  
Jul-Aug;4(4):231-43.

## Manufacturer Information

---

Abacavir sulfate, Lamivudine, and Zidovudine  
GlaxoSmithKline  
5 Moore Drive  
Research Triangle Park, NC 27709  
(888) 825-5249

## For More Information

---

Contact your doctor or an AIDSinfo Health  
Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday,  
12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: [http://aidsinfo.nih.gov/live\\_help](http://aidsinfo.nih.gov/live_help)  
Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

# Abacavir sulfate, Lamivudine, and Zidovudine



## References

---

1. GlaxoSmithKline - Trizivir Product Information, August 2002; p. 1. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
2. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 10, 24. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
3. Protocol ID: ACTG A5095 -
4. NIH/NIAID - Press Release; Interim Results from AACTG Protocol A5095. Available at: [http://www.aidsinfo.nih.gov/document/data/aprs\\_lib/aprs\\_A00659.html](http://www.aidsinfo.nih.gov/document/data/aprs_lib/aprs_A00659.html). Accessed 1/6/04.
5. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 3-5. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
6. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 3-6. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
7. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 3-6. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
8. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 5. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
9. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 7, 18-19. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
10. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 4-5. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
11. NIH/NIAID - Press Release; Interim Results from AACTG Protocol A5095. Available at: [http://www.aidsinfo.nih.gov/document/data/aprs\\_lib/aprs\\_A00659.html](http://www.aidsinfo.nih.gov/document/data/aprs_lib/aprs_A00659.html). Accessed 1/6/04.
12. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 20-21. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
13. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 14. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
14. USP DI - 2003; p. 2791.
15. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 14. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
16. AHFS Drug Information - 2003; p. 668.
17. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 14. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
18. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 23. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
19. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 6. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
20. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 7-8. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
21. USP DI - 2003; p. 3.
22. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 16. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
23. USP DI - 2003; pp. 1647-48.
24. USP DI - 2003; pp. 2789-90.
25. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 9. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
26. USP DI - 2003; pp. 2789-90.
27. GlaxoSmithKline - Trizivir Product Information, August 2002; p. 16. Available from: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
28. USP DI - 2003; p. 2789.

# ***Abacavir sulfate, Lamivudine, and Zidovudine***



29. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 16. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
30. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; pp. 6-7. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
31. USP DI - 2003; p. 4.
32. GlaxoSmithKline - GlaxoSmithKline, Trizivir Product Information, August 2002; p. 24. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
33. GlaxoSmithKline - Trizivir Product Information, August 2002; p. 24. Available from: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
34. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 17.
35. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 477.
36. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 920.
37. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 17.
38. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 477.
39. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 920.
40. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 12/22/03.
41. Merck Index - 2001; pp. 1, 958, 1809.
42. Merck Index - 2001; pp. 1, 958, 1809.
43. Merck Index - 2001; 13th ed; pp. 1, 958, 1809.
44. GlaxoSmithKline - Trizivir Product Information, August 2002; p. 1. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.
45. GlaxoSmithKline - Trizivir Product Information, August 2002; pp. 2-3. Available at: [http://us.gsk.com/products/assets/us\\_trizivir.pdf](http://us.gsk.com/products/assets/us_trizivir.pdf). Accessed 12/29/03.